EXTREMAL GRAPH THEORY VIA TWO GEOMETRY QUESTIONS

Haydar Göral

Department of Mathematics, Izmir Institute of Technology, Urla, Izmir, Turkey haydargoral@iyte.edu.tr

Abstract

In this short note, we will give the basic results of extremal graph theory motivated by the two geometry problems.

1. Two Geometry Questions

Motivating questions related to geometry:

- 1. Given n points in the plane (\mathbb{R}^2), at most how many of them can be unit distance apart? (Erdős Unit Distance Problem, 1946)
- 2. Let $n_0 \geq 2$ be given. Find n such that, any n point in \mathbb{R}^2 , no three of them are collinear, contains n_0 points in *convex position* (all points are need for the convex hull).

2. Notations

G = (V, E): Graph with vertex set V and edge set E (simple graph)

 K_m : The complete graph on m vertices

 $K_{s,t}$: The complete bipartite graph (V_1, V_2, E) where $|V_1| = s, |V_2| = t$

 K_{s_1,\ldots,s_r} : The complete r-partite graph (V_1,\ldots,V_r,E) where $|V_i|=s_i$

deg(v): The degree of the vertex v

e(G): The cardinality of edge set E, where G = (V, E)

R(n,k): Ramsey Number

ex(n, H): The extremal number of H

 $\chi(H)$: The chromatic number of H

 $T_{n,r}$: Turán Graph

e(X,Y): The number of edges between X and Y

d(X,Y): The density of X and Y

 \mathbb{P} : The set of all prime numbers

[n]: The set of first n positive integers

 $\mathbb{Z}_{>0}$: The set of non-negative integers

3. Ramsey Numbers

R(n,k): The minimal number m such that for every 2-coloring of K_m $\left(K_m:\binom{[m]}{2}\right)=\{A\subseteq [m]:|A|=2\}\right)$ there exists a blue $K_n\subseteq K_m$ or there exists a red $K_k\subseteq K_m$ i.e. there exists a subset $X\subseteq [m]$ with n elements and $\binom{X}{2}$ (subsets of X of size 2) is colored by blue, or there exists a subset $Y\subseteq [m]$ with k elements and $\binom{Y}{2}$ is colored by red.

- Observe that R(n,2) = n, R(2,k) = k.
- For n, k > 2, one can show that

$$R(n,k) \le R(n-1,k) + R(n,k-1).$$

By induction, this proves the existence of the Ramsey numbers.

- Graph: 2-graph Hypergraph: s-graph (an edge is a set containing s elements!)
- $R_s(n,k)$, $(s \ge 2)$: For s-graphs, we color subsets of size s.
- We have the following bound $R_s(n,k) \le 1 + R_{s-1}(R_s(n-1,k), R_s(n,k-1))$. Also, $R_s(n,s) = n$.
- Instead of 2 colors, we can use finitely many colors:

$$R(n_1,\ldots,n_k) \leq R(n_1,\ldots,n_{k-2},R(n_{k-1},n_k)),$$

(existence of K_{n_i} of color i for some i). Similarly, one has

$$R_s(n_1,\ldots,n_k) \leq R_s(n_1,\ldots,n_{k-2},R_s(n_{k-1},n_k)).$$

This proves the existence of the Ramsey numbers (multi versions).

4. Forbidding a Subgraph

Theorem 1. (Mantel, 1907) Every graph on n vertices with edge density greater than $\frac{1}{2} \cdot \frac{n}{n-1}$ contains a triangle.

Theorem 2. (Roth, 1953) Every subset of $\mathbb{Z}_{\geq 0}$ with positive upper density contains a 3-term arithmetic progression (3-AP).

We also note the following two far-reaching results of Szemerédi and Green-Tao.

- Szemerédi (1975): If $A \subseteq \mathbb{Z}_{\geq 0}$ and $\bar{d}(A) > 0$, then A contains a k-AP for every $k \geq 3$.
- Green-Tao (2005): P contains arbitrarily long APs.

Mantel's theorem is equivalent of the following:

Theorem 3. (Mantel) Any triangle-free graph on n vertices has at most $\lfloor \frac{n^2}{4} \rfloor$ edges.

Proof. Let $A \subseteq V$ be a maximum independent set (no edges in A). Observe that the neighborhood of any vertex v is independent by triangle-freeness:



So, $\deg(v) \leq |A|$. Set $B = V \setminus A$. Then, every edge must have a vertex in B. Therefore,

$$e(G) \le \sum_{v \in B} \deg(v) \le |A||B| \le \left(\frac{|A| + |B|}{2}\right)^2 = \frac{n^2}{4}.$$

Remark 1. For equality, we need |A| = |B|, no edge in B as well, $\deg(v) = |A|$ for all $v \in B$:

 $K_{\frac{n}{2},\frac{n}{2}}$ for n even or $K_{\frac{n-1}{2},\frac{n+1}{2}}$ for n odd.

Turán Graph:


 $T_{n,r}$ is the complete r-partite graph $K_{s_1,...s_r}$ with

$$n = s_1 + \dots + s_r$$
 and $s_i \in \left\{ \left\lfloor \frac{n}{r} \right\rfloor, \left\lceil \frac{n}{r} \right\rceil \right\}$.

Example: $T_{8,3} = K_{2,3,3}$ (edges only between V_i, V_j for $i \neq j$)




• $e(T_{n,r}) \le (1 - \frac{1}{r}) \frac{n^2}{2} = \binom{r}{2} \frac{n}{r} \cdot \frac{n}{r}$. One can easily see this when $r \mid n$.



Theorem 4. (Turán) Any K_{r+1} -free graph on n vertices has at most $e(T_{n,r})$ edges. $(T_{n,r})$ is the unique maximizer)

Proof. We proceed by induction on n. If $n \leq r$, then $T_{n,r} = K_n$, and it is the best option. Suppose n > r and the statement holds for all graphs with less than n vertices. Let G be a K_{r+1} -free graph with n vertices and it has the maximal number of edges. Thus, G has a copy of K_r , otherwise we add an edge. Let A be a set of vertices which form a K_r , and set $B = V \setminus A$.

For every $v \in B$, it has at most r-1 neighbors in A.



Hence,

$$e(G) \le {r \choose 2} + (r-1)(n-r) + e(T_{n-r,r}) = e(T_{n,r}).$$

• ex(n, H) = the maximum possible number of edges in a H-free graph on n vertices: the extremal number of H

• $\chi(H) = the chromatic number of H$

Examples:

$$\chi(K_{r+1}) = r + 1$$
$$\chi(T_{n,r}) = r$$

$$ex(n, K_{r+1}) = e(T_{n,r}) = \left(1 - \frac{1}{r} + o(1)\right) \binom{n}{2}$$

If $\chi(H) = r + 1$, then $T_{n,r}$ is indeed H-free and

$$ex(n, H) \ge ex(T_{n,r}) = \left(1 - \frac{1}{r} + o(1)\right) \binom{n}{2}.$$

How far can we go? This can be answered by the following result of Erdős-Stone-Simonovits.

Theorem 5. (Erdős-Stone-Simonovits, 1946)

$$ex(n,H) = \left(1 - \frac{1}{\chi(H) - 1} + o(1)\right) \binom{n}{2}$$

• Petersen graph = H, $\chi(H) = 3 = \chi(K_3)$



• There is a proof of ESS theorem which applies Szemerédi's Regularity Lemma. Now, we recall Szemerédi's Regularity Lemma briefly.

For $X, Y \subseteq V$, define

$$e(X,Y) = |\{(x,y) \in X \times Y : xy \in E(G)\}|$$

$$d(X,Y) = \frac{e(X,Y)}{|X||Y|}.$$

 ε -regular pair: Let G be a graph and $U, W \subseteq V$. We call (U, W) an ε -regular pair in G if for all $A \subseteq U, B \subseteq W$ with $|A| \ge \varepsilon |U|, |B| \ge \varepsilon |W|$, one has:

$$|d(U, W) - d(A, B)| \le \varepsilon.$$

 ε -regular partition: The partition $P = \{V_1, \dots V_k\}, V_1 \sqcup \dots \sqcup V_k = V$, is an ε -regular partition if

$$\sum_{\substack{(i,j)\in[k]^2\\(V_i,V_j)\text{ not }\varepsilon-\text{regular}}} |V_i||V_j|\leq \varepsilon |V|^2.$$

Lemma 1. (Szemerédi's Regularity Lemma (SGL)) For every $\varepsilon > 0$, there exists $M \in \mathbb{Z}_{\geq 0}$ such that every graph has an ε -regular partition into at most M parts.

We know that SGL \implies Roth's Theorem, SGL \implies ESS and SGL \implies Szemerédi's Theorem (with more arguments), and hyper-regularity \implies multi-dimensional Szemerédi's Theorem.

• Zarankiewicz Problem: s < t

Theorem 6. (Kővári–Sós–Turán, 1954)

$$ex(n, K_{s,t}) = O_{s,t}(n^{2-\frac{1}{s}})$$

Conjecture 1. (KST)

$$ex(n, K_{s,t}) \ge c_{s,t} n^{2-\frac{1}{s}}$$

ESS open for hypergraphs: Even for $K_4^{(3)}$ (bounds via flag algebras)

Back to Erdős unit distance problem: Q1

Given n points in the plane, at most $O(n^{\frac{3}{2}})$ pairs are unit distance apart.

Proof. Consider the graph on those points with edges indicating unit distance. This graph is $K_{2,3}$ -free as circles only intersect twice. We are done by KST.



Conjecture 2. The true bound for the Erdős unit distance problem is $O_{\varepsilon}(n^{1+\varepsilon})$.

Theorem 7. (Erdős-Rényi-Sós, 1966)

$$ex(n, K_{2,2}) \ge \left(\frac{1}{2} - o(1)\right) n^{\frac{3}{2}}$$

Proof. Let $p=(1-o(1))\sqrt{n}$ be a prime in $[x-x^{0.525},x]$ for $x=\sqrt{n}$, by Baker-Harman-Pintz, 2007.

Consider G = (V, E) as follows:

for
$$a, b = 0, 1, \dots, p - 1$$
 , $(a, b) \neq (0, 0)$


$$x, y = 0, 1, \dots, p - 1$$
 , $(x, y) \neq (0, 0)$

there is an edge between (a,b) and (x,y) if $ax+by\equiv 1\pmod p$. (There are p^2-1 many such pairs.) For other $n-p^2+1$ vertices, they have degree =0, i.e. isolated. Notice that

$$ax + by \equiv 1 \pmod{p}$$
,

$$cx + dy \equiv 1 \pmod{p}$$

has at most one solution.



So G is $K_{2,2}$ —free, and

$$e(G) = \frac{1}{2} \sum_{v \in V} \deg(v) \ge \frac{1}{2} (p^2 - 1)(p - 1) = \left(\frac{1}{2} - o(1)\right) n^{\frac{3}{2}}.$$

(a line ax + by = 1 has at least p - 1 elements)


• Brown (1966): $ex(n, K_{3,3}) \ge \left(\frac{1}{2} - o(1)\right) n^{\frac{5}{3}}$

• KST for $K_{4,4}$ open!

Back to Q2:


Theorem 8. (Erdős-Szekeres, 1935) Let $n_0 \geq 2$. There exists a positive integer n such that: Any set of n points in \mathbb{R}^2 , no three of which are collinear, contains n_0 points in convex position.

Proof. Let $n = R_4(n_0, 5)$. We give a 4-subset color blue if they are in convex position, otherwise the color is red.



Ramsey theorem ensures the existence of an n_0 -subset $S \subseteq [n]$ with all 4-subsets are colored by blue, or a 5-subset with all 4-subsets are colored by red.

First Case: This S is in convex position. Suppose some points are not needed and consider a minimal subset which gives our convex hull (a polygon):




We obtain a 4-subset which is not in convex position. So, we are done.

Second Case: This is impossible due to the $Happy\ Ending\ Problem$ by Esther. \Box

Lemma 2. (Happy Ending Problem) Every set with 5 points in \mathbb{R}^2 , no three of them lie on a line, always contains 4 points in convex position.

Proof. The following figure is the proof:



Enough to check: 2,3,4,5 (BCDE), 14 (ABDE), 15 (ACDE)

Acknowledgement: This note was made possible thanks to Doruk Üstündağ's keen attention during the lecture and his efforts in typing it up.

References: Yufei Zhao, Graph theory and additive combinatorics. Cambridge University Press (2021).